

Vaccination protocols:

A brief description of the below diseases follow the protocol tables:

Disease	Vaccination protocols for Adult Horses			
	Previously vaccinated	Naïve/unvaccinated		Broodmares
Tetanus*	Every 12 months	Every 12 months	Boost 3-6 weeks after initial vaccine	1 month prior to foaling
EEE/WEE*	Every 12 months	Every 12 months	Boost 3-6 weeks after initial vaccine	1 month prior to foaling
WNV* (West Nile Virus)	Every 12 months -Spring-summer -*6months with high exposure	Every 12 months Spring-summer -*6 months with high exposure	Boost 3-6 weeks after initial vaccine	1 month prior to foaling
EHV 1/4 Aka: Rhino	Every 6-12 months depending on exposure	Every 6-12 months depending on exposure	Boost 3-6 weeks after initial vaccine	At 5, 7, 9 months of pregnancy. -Abortion prevention
EIV (Influenza)	Every 6-12 months depending on exposure	Every 6-12 months depending on exposure	Boost 3-6 weeks after initial vaccine	1 month prior to foaling
Strep Equi (Strangles)	Every 12 months	Every 12 months	Boost 3-6 weeks after initial vaccine	

Vaccination Protocols for Foals				
Disease	Vaccinated Mares		Unvaccinated Mares	
	6 months of age	7 months of age	3-4 months of age	4-5 months of age
Tetanus	√	√	√	√
EEE/WEEE	√	√	√	√
WNV	√ -May depend on season	√ -May depend on season	√ -May depend on season	√ -May depend on season
EHV ¼	√	√ -booster 6-12 months depending on exposure	√	√ -Booster 6-12 months depending on exposure
EIV	√	√	√	√
Strep Equi	Depending on exposure		Depending on exposure	